

DATA SHEET



AKIPOR 6.3®

Presentation

Freeze-dried pellet and oily diluent for injectable emulsion.
Each 2-ml dose of vaccine contains at least $10^{6.3}$ CCID₅₀ of Live Aujeszky's disease (syn: pseudorabies) virus, gE-deleted Bartha strain. In addition, AKIPOR 6.3 contains an oily diluent containing 0.2 mg of Thiomersal, at most and 2 ml of oily excipient.

Indication

Active immunization against Aujeszky's disease and passive immunization of piglets born to vaccinated sows via colostral antibodies.

Dosage and administration

A 2 ml dose injected intramuscularly into the neck in the area behind the ear, according to the following schedule:

In breeder pigs

Primary vaccination

Replacement gilts: 2 injections at a 3-4 week interval before the first service
Active breeding sows: 2 injections at a 3-4 week interval, regardless of gestational status

Booster injections

1 injection 2-3 weeks before each subsequent farrowing.

In fattening pigs

1 injection at the beginning of the fattening period, at the minimum age of 10 weeks
In contaminated farms, perform a 2nd injection 4 weeks later

Contra-indications, warnings, etc

There are no contra-indications.

Do not vaccinate species other than pigs

Vaccinate only healthy animals.

Apply usual procedures for the handling of the animals.

Shake before use.

Use immediately after reconstitution of the freeze-dried pellet with the oily diluent.

Use sterile equipment free from antiseptic and/or disinfectant for vaccine solution preparation and injection purposes.

If the vaccine is accidentally injected to man, urgent medical attention is necessary.

Apply usual aseptic procedures.

Pharmaceutical precautions

Store between +2°C and +8°C, protected from light.

Further information

Vaccination during pregnancy or lactation is not contraindicated.

No undesirable effect has been observed after the administration of an overdose of vaccine.

Merial Asia Pte Ltd
57A Science Park Drive,
The Faraday
Singapore Science Park 1
Singapore 118238